

OPERATIONS AND IDENTITIES IN TENSOR ALGEBRA

Chernyakov Yu. [†], Dolotin V. [‡]

Abstract

We define the class of non-decomposable N -ary operations in the mixed tensor algebra $\bigoplus_{i,j=0}^{\infty} A_i^j$. There are higher Jacobi-like identities for (binary) deformed matrix commutator and a 3-ary operation which is non-decomposable into binary ones.

1 Introduction

The subject of the algebra is mostly deals with studying the action of the rings of linear maps on the corresponding modules ([1],[2]). Superposition of these maps stipulates the binarity of the composition law in the space of operators. Linear combinations of such superpositions give new (generically, non-associative) composition laws, such as multiplications in Lie algebra, Jordan algebra and their deformations. Let tensor algebra be generated by a given linear space V . From the tensor algebra point of view the operator ring is the space $V \otimes V^*$ of (1,1)-type tensors. The composition law in such space is generated by elementary operations for generators:

$$(e_i \otimes e_j^*) \cdot (e_k \otimes e_l^*) = (e_j^*, e_k) e_i \otimes e_l^* = \delta_{jk} e_i \otimes e_l^*$$

However, the class of such elementary operations in the mixed tensor algebra $A_{\bullet}^{\bullet} := \bigoplus_{i,j=0}^{\infty} A_i^j = \bigoplus_{i,j=0}^{\infty} V^{\otimes i} \otimes V^{*\otimes j}$ is considerably wider. Combinations of these elementary operations generate the class of operations which in general is non-binary and is not decomposable into binary operations. So, there appears a problem of finding such subspaces in $A_{\bullet}^{\bullet}$, which are invariant with respect to these operations, and finding the identities which these operations satisfy.

This paper is organized in the following way. In Chapter 2 we describe the language which simplifies the further consideration. In Chapter 3 the finite dimensional subalgebras of mixed tensor algebra $\bigoplus_{i,j=0}^{\infty} A_i^j$, i.e. the (1,1)-type tensors (matrices), binary operations and the corresponding identities are considered. In Chapter 4 we consider the tensor space $A_2^1 \oplus A_1^2$, a 3-ary

[†]ITEP, Moscow, Russia, e-mail: chernyakov@itep.ru

[‡]ITEP, Moscow, Russia, e-mail: vd@itep.ru

operation non-decomposable into binary operations and an identity which generalizes the commutator and Jacobi identity. In Chapter 5 we speak about the n -ary operations. The Chapter 6 is a discussion.

2 Language

Let us consider the action of the set of linear operators on the vector space V . The dual space V^* consists of linear forms on V . It is possible to define the operation called contraction, given the value of each element of V^* on the elements of V . The result of this operation is scalar. It is possible to define another operation called tensor product of V^* and V . The linear operator is defined as (1,1)-type tensor where digits denote the number of "upper and lower indices" of the tensor correspondingly, i.e. $A_1^1 = V \otimes V^*$. The spaces V^* and V generate the tensor algebra. We consider it as a graded space. The value of this grading is defined as the difference between the number of lower and upper indexes of the tensor. Thus, the grading of V is $[V] = +1$ and of V^* is $[V^*] = -1$.

It is convenient to describe the general situation by the following picture:

$$\dots \oplus \begin{bmatrix} A_1^0 \\ \oplus \\ A_2^1 \\ \oplus \\ A_3^2 \\ \oplus \\ A_4^3 \\ \dots \end{bmatrix} \oplus \begin{bmatrix} A_0^0 \\ \oplus \\ A_1^1 \\ \oplus \\ A_2^2 \\ \oplus \\ A_3^3 \\ \dots \end{bmatrix} \oplus \begin{bmatrix} A_0^1 \\ \oplus \\ A_1^2 \\ \oplus \\ A_2^3 \\ \oplus \\ A_3^4 \\ \dots \end{bmatrix} \oplus \dots \quad (1)$$

Let us introduce some diagrammatic language which helps us to produce further results. We use the notation $\xrightarrow{a}$ for the (1,1)-type tensor where the arrow tip means the upper index and the tip without the arrow means the lower index of the tensor. Using this rule we get for the tensors $a \in A_3^1$ and $b \in A_2^5$ correspondingly (Fig.1).

Following this notation the contraction of two (1,1)-type tensors A and B using the upper index of A and the lower of B looks as follows figure 2.

As a result we get a (1,1)-type tensor again. Here and further we take into account that all indices have equal number of values. So, the contraction between arbitrary two indexes of different types (upper and lower) is possible. For example, the tensor product $\otimes$ is represented as figure 3.

In this case we get (2,2)-type tensor. Note, that similar notations were used in the paper [3], but being applied to differential operators, which may be regarded a the subclass of symmetric tensors in $A_\bullet^\bullet$.

In the mixed tensor algebra $A_\bullet^\bullet$ there exist subspaces which are invariant with respect to some fixed operations (non necessarily binary). These subspaces may be called subalgebras of $A_\bullet^\bullet$. In turn, these operations may satisfy certain identities.




Figure 1:



Figure 2:




Figure 3:




Figure 4:

3 (1,1)-type tensors - matrices and binary operations

Let us consider the following question: how many different types of contractions (called further elementary operation) of two (1,1)-type tensors is there. We consider as the result of the contraction tensors having the same type as the type of ingredients and tensors of other types as well. One can check that using the contraction and tensor product (which corresponds to the contraction of empty set of indices) we can get seven variants of difference compositions (Fig.4).

The first four variants are the (1,1)-type tensors, the next two ones are the (0,0)-type tensors - scalars, and the latter one is (2,2)-type tensor. Note that the result of each elementary operation has the grading equal to zero. Evidently this is the property of contraction only for (1,1)-type tensors, or other zero graded $((m,m)$ -type) tensors.

Further in this Chapter we will be interested in such elementary operations having (1,1)-type tensors as the result. These operations transfer the initial space, the space of (1,1)-type tensors, to itself. Now let us define a generic binary operation as a linear composition of elementary operations between tensors A and B (here we do not write tensor indices applied to (1,1)-type tensors):

$$A \circ B = \alpha \cdot AB + \beta \cdot BA + \gamma \cdot A \operatorname{Tr} B + \delta \cdot B \operatorname{Tr} A \quad (2)$$

where $\alpha, \beta, \gamma, \delta \in C$. So, we define the binary bilinear operation φ_2 on the matrix space:

$$\varphi_2 : A_1^1 \times A_1^1 \rightarrow A_1^1 \quad (3)$$

Note, if we put $\alpha + \beta = 0$, $\gamma = 0$, $\delta = 0$, we get

$$A \circ B = \alpha \cdot (AB - BA), \quad (4)$$

It is the matrix commutator if we put $\alpha = +1$, $\beta = -1$. How can we see the Jacobi identity using the language defined in Chapter 2 ? We have to consider the three-symbol words, for example $a b c$ where each symbol corresponds to the (1,1)-type tensor and each word is the result of contractions of these tensors (Fig.5).




Figure 5:

It is possible to get this word by two ways using the primary operations: $A \circ B$ and $B \circ C$. On the second step of this procedure we must operate with the result of the first primary operation and the rest third tensor. Organizing the enumeration in the appropriate way (using the permutations) we get the Jacobi identity:

$$(A \circ B) \circ C + (B \circ C) \circ A + (C \circ A) \circ B = 0. \quad (5)$$

For the operation (2) it is possible to get the identity setting $\beta = -\alpha$, $\delta = -\gamma$. This identity is the generalization of the Jacobi identity (see details in Appendix 1):

$$\begin{aligned} & ((A \circ B) \circ C) \circ D + ((C \circ B) \circ D) \circ A + ((C \circ D) \circ A) \circ B + ((A \circ D) \circ B) \circ C + \\ & ((C \circ A) \circ B) \circ D + ((D \circ C) \circ B) \circ A + ((A \circ C) \circ D) \circ B + ((B \circ A) \circ D) \circ C + \\ & ((B \circ C) \circ A) \circ D + ((B \circ D) \circ C) \circ A + ((D \circ A) \circ C) \circ B + ((D \circ B) \circ A) \circ C \\ & = 0. \end{aligned} \quad (6)$$

In order to get this identity we use the following arguments. First of all, the direct testing shows that ordinary cyclic permutations of symbols in $(A \circ B) \circ C$ does not give zero result, so we can take triple compositions of the operation (2). In this case we must take into account all primary operations taking part in formation of four-symbolic words. There are twelve such primary operations: $(AB), (BA), (CA), (AC), (AD), (DA), (BC), (CB), (BD), (DB), (DC), (CD)$. It is possible to rewrite the identity (6) in more adequate and general form. For a 4-tuple of elements A_1, A_2, A_3, A_4 we get the following trilinear operation φ_3 :

$$\begin{aligned} & \varphi_3(A_1, A_2, A_3) = \\ & \sum_{i=1}^3 (-1)^{i-1} \varphi_2(\varphi_2(\overline{A_i}), A_i) = \{ \{A_2, A_3\}, A_1 \} - \{ \{A_1, A_3\}, A_2 \} + \{ \{A_1, A_2\}, A_3 \} \end{aligned} \quad (7)$$

where $\overline{A_i}$ denotes the complement to A_i in the triple A_1, A_2, A_3 , φ_2 is the primary binary operation (2). Using φ_2 and φ_3 , we get:

$$\varphi_4(A_1, A_2, A_3, A_4) = \sum_{i=1}^4 (-1)^{i-1} \varphi_2(\varphi_3(\overline{A_i}), A_i) \quad (8)$$

And for φ_2 with $\alpha + \beta = 0$, $\gamma + \delta = 0$ we get

$$\varphi_4(A_1, A_2, A_3, A_4) \equiv 0. \quad (9)$$

Let us formulate the following

Proposition 1: *For the operation (2) with $\alpha + \beta = 0$, $\gamma + \delta = 0$ it is possible to define the combinatoric identity (9) using only the operations (2).*

Note that the identity (9) uses higher operations obtained from the binary operations (2). This is similar to the existence of Jacobi identity in matrix algebra defined through the commutation operation.




Figure 6:

4 The tensor space $\mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ and 3-ary operation

Now let us take a pair of tensors A, B in general having a non-zero gradings $[A] = n, [B] = k$. For any type of contractions the result is always in the space with grading $n + k$. The naturally arising question is what can be done in order to remain in initial grading? The answer is that if we take a space of linear combination of tensors having different gradings there are operations with the result being tensor from this original space, so the grading type is preserved.

Let us make the following observation. For elements $A \in \mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ of the direct sum of pure graded spaces it is possible to find an operation keeping the result in initial space. To do this we take two more such linear combinations $B, C \in \mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ and consider various collections of triples of tensors, each being a pure graded component in the sums $A = A_2^1 + A_1^2, B = B_2^1 + B_1^2$ and $C = C_2^1 + C_1^2$, with a set of appropriate contractions. We exclude the contractions of a tensor with itself. These conditions define the following rule: in order to get a tensor from the space $\mathcal{A}_1^2$ one need to take two (2,1)-type tensors and one (1,2)-type tensor and for getting a tensor from the space $\mathcal{A}_2^1$ one need to take two (1,2)-type tensors and one (2,1)-type tensor. The number of elementary operations is fourteen: seven for each type of resulting tensors (Fig.6 and Fig.7).

We see that in contrast to the matrix multiplication in the space $\mathcal{A}_1^1$ in the case of tensor space $\mathcal{A} = \mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ the important role plays the type of contractions, in other words the topology of the corresponding diagram. The diagrams of the first two elementary operations may be aligned (schematically) in a line because the objects in this case are placed one after another. We called these operations by operations of "linear" type. The diagrams of the other elementary operations can be placed on the plane.

Note one important property of the operation on the space $\mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ - they are non-decomposable into binary operations because the result of any binary operation may not belongs to the initial space $\mathcal{A}_2^1 \oplus \mathcal{A}_1^2$. The origin of these 3-ary operations may be traced from the rule of the action of $\mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ on the composed "vector space" $V^{\otimes 2} \oplus V^{\otimes 1} = \mathcal{A}^2 \oplus \mathcal{A}^1$, which is




Figure 7:

analogous to the linear representation for matrix algebra:

$$(A_2^1 + A_1^2)(v^2 + v^1) = A_2^1 \cdot v^2 + A_1^2 \cdot v^1$$

So under this action the space $V^{\otimes 2} \oplus V^{\otimes 1}$ passes to itself.

So, taking a linear combination of the elementary operations we get a generic 3-ary operation:

$$\varphi : \mathcal{A} \times \mathcal{A} \times \mathcal{A} \longrightarrow \mathcal{A}, \quad (10)$$

while for any binary operation

$$\tau(\mathcal{A} \times \mathcal{A}) \not\subset \mathcal{A}. \quad (11)$$

Now let taking a linear combination of two first elementary operations we define a 3-ary operation and a Jacobi-like identity. Let us define the 3-commutator with the condition $\alpha + \beta + \gamma = 0$, which is analogous to the condition $\alpha + \beta = 0$ for the commutator (Fig.8):

$$\begin{aligned} (X, Y, Z) = & \\ & \alpha \cdot X_1^2 Y_2^1 Z_1^2 + \alpha \cdot Z_1^2 Y_2^1 X_1^2 + \beta \cdot Z_1^2 X_2^1 Y_1^2 + \beta \cdot Y_1^2 X_2^1 Z_1^2 + \gamma \cdot X_1^2 Z_2^1 Y_1^2 + \gamma \cdot Y_1^2 Z_2^1 X_1^2 + \\ & + \alpha \cdot X_2^1 Y_1^2 Z_2^1 + \alpha \cdot Z_2^1 Y_1^2 X_2^1 + \beta \cdot Z_2^1 X_1^2 Y_2^1 + \beta \cdot Y_2^1 X_1^2 Z_2^1 + \gamma \cdot X_2^1 Z_1^2 Y_2^1 + \gamma \cdot Y_2^1 Z_1^2 X_2^1, \end{aligned} \quad (12)$$

where:

$$X = X_2^1 + X_1^2, \quad Y = Y_2^1 + Y_1^2, \quad Z = Z_2^1 + Z_1^2, \quad (13)$$

Here is assumed the summation of the different combinations of tensor contractions. So, the number of terms in the 3-commutator is 24. But we can insert the 3-commutator under the sign of summation and in further we do not specially stipulate the appeared adding terms from the summing the different tensor contractions. We set α, β, γ equal to the different values of the cubic root on unity:

$$\alpha = 1, \quad \beta = e^{i\frac{2\pi}{3}}, \quad \gamma = e^{i\frac{4\pi}{3}}. \quad (14)$$




Figure 8:

Thus we can consider 3-commutator modulo any number α, β, γ . This is analogue of $\alpha = -\beta = 1$. By analogy to the antisymmetry of the commutator:

$$[X, Y] + [Y, X] = 0, \quad (15)$$

the 3-commutator has the following property:

$$(X, Y, Z) + (Z, X, Y) + (Y, Z, X) = 0, \quad (16)$$

As the result of n -times action of the 3-commutator on the tensor space $\mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ we get the linear combination of the different words of $2n + 1$ symbols. After the second action of the 3-commutator there appear words consisting of five symbols, like $A_1^2 B_2^1 C_1^2 D_2^1 E_1^2$, $A_2^1 B_1^2 C_2^1 D_1^2 E_2^1$ and so on. Using the 3-commutator it is possible to define the identity which consist of all such words. Let us take as initial the word $A_1^2 B_2^1 C_1^2 D_2^1 E_1^2$. Note that this word can be obtained from the three different primary 3-commutator operations:

$$(A, B, C), \quad (B, C, D), \quad (C, D, E). \quad (17)$$

It is possible to get this operation by the cyclic permutation in the primary operation (A, B, C) , and the secondary one $((A, B, C)D, E)$. In this case the initial word is $A_1^2 B_2^1 C_1^2 D_2^1 E_1^2$. Thus we set the order of writing the identity although we can take any operation as a primary one. In a similar way, dealing with the Jacobi identity we take as initial the word ABC and the operation $[A, B]$ as primary, and then make cyclic permutations. One needs to take all primary 3-symbol operations. In other words, all sets which consist of three symbols chosen from the given five ones. So we have 10 primary operations modulo the coefficients (weights). It is possible to get all these operations from the two 5-symbol classes: $((A, B, C)D, E)$ and $((D, A, C)E, B)$. We must also take into account the existence of classes which differ by cyclic permutations in the sets of three symbols: $(X, Y, Z) + (Z, X, Y) + (Y, Z, X) = 0$ - $(Z, Y, X) + (X, Z, Y) + (Y, X, Z) = 0$. For a chosen initial word it does not matter which operation we define as the primary one: (A, B, C) or (C, B, A) . If we start from (C, B, A) , it is possible to get the same sets of tree symbol as for another four primary operations (as in the case of (A, B, C)), but belonging to

another class. Note that in the case of usual Jacobi identity we have only cyclic permutations between three symbols because the primary operations consist of two symbols and belong to the unique class of cyclic permutations. The identity must have symmetry with respect to the different classes. The identity is expressed in the following form:

$$\begin{aligned}
& ((A, B, C)D, E) + ((B, C, D)E, A) + ((C, D, E)A, B) + ((D, E, A)B, C) + ((E, A, B)C, D) + \\
& + ((C, B, A)E, D) + ((B, A, E)D, C) + ((A, E, D)C, B) + ((E, D, C)B, A) + ((D, C, B)A, E) + \\
& + ((D, A, C)E, B) + ((A, C, E)B, D) + ((C, E, B)D, A) + ((E, B, D)A, C) + ((B, D, A)C, E) + \\
& + ((C, A, D)B, E) + ((A, D, B)E, C) + ((D, B, E)C, A) + ((B, E, C)A, D) + ((E, C, A)D, B) + \\
& = 0
\end{aligned} \tag{18}$$

For more detail about this identity see Appendix 2. Let us formulate the following

Proposition 2: *For the 3-ary operation (12) non-decomposable into binary ones there is a combinatorial identity (18).*

This is similar to the existence for the matrix commutator Jacobi identity defined by the commutation operation.

5 N-ary operation

There are many various n-ary operations in the mixed tensor algebra $\bigoplus_{i,j=0}^{\infty} A_i^j$. For example, there are 3-ary operation for $A_1^2 \oplus A_4^3$ with the five contractions, 5-ary for $A_1^2 \oplus A_5^2$ with the eight contractions and so on, and, possibly, the different identities. The meaning of these identities as analogues of Jacobi identity has the combinatorial nature. It pertains to the possibility to come to the same contraction diagram by different pathes having different weights. Under the "path" we mean the following: the word can be obtained by the different primary and secondary operations, and the sequence of this operations is called the path.

Being restricted to the case of "linear" elementary operations for some operator spaces let us consider the following table:

Operators	Space	Rank of the elementary operation
A_1^1	V^1	2
$A_1^2 \oplus A_2^1$	$V^1 \oplus V^2$	3
$A_1^3 \oplus A_2^1 \oplus A_3^2$	$V^1 \oplus V^2 \oplus V^3$	4
...	...	...
$A_1^n \oplus A_2^1 \oplus A_3^2 \oplus \dots \oplus A_n^{n-1}$	$V^1 \oplus V^2 \oplus \dots \oplus V^n$	$n + 1$

(19)

In the column "Space" there are examples of the vector spaces on which the tensors from column "Operator" act as operators.

Although the existence of the different generalized identities and corresponding operations (composed by n-ary elementary operations) is the subject of the further research let us define some basis principles of construction the operations and identities. First of all we give some remarks about the 3-ary operation (12). Defining this operation we chose the second position in the sequence of the objects A_1^2 and A_2^1 and assumed the following: when Y has the second

position then the weight of the elementary operation is α , when $X - \beta$, and when $Z - \gamma$. Note that we can chose the first and the third position.

For example, let us consider the operators of the space $A_1^3 \oplus A_2^1 \oplus A_3^2$ and use the principle stated above. In this case there is the primary operation consisted of four symbols $(XYZW)$. After twice acting of this operation we get the seven-symbolic words. Then the number of such operations is 35. Choosing the first position we assume the following: when X has the first position then the weight of the elementary operation is α , when $Y - \beta$, when $Z - \gamma$ and when $W - \gamma$. We get that the number of the elementary operations with the equal weight is 18 for the all type of the operators. In the other hand the number of the elementary operations with the different weight is 6 for the each type of the operators. The total number of the elementary operations in each primary operation is 72. In the identity the number of terms must be $35 \cdot 2 = 72$. They are divided in groups having 7 terms in according to the number of terms in each class of the cyclic permutations. Each word must occur 8 times. We suppose that this principles can be used in the cases of any n -ary operation.

6 Discussion

We suppose that the following questions must be interesting. First of all there are general questions about operations. Given an operation in $\mathcal{A}_\bullet$: what is the set of corresponding identities (in particular, for operations on $\mathcal{A}_2^1 \oplus \mathcal{A}_1^2$ having diagrams with "non-linear" topology)? When can we restrict to double iterations of the primary operation in the identity?

The operation (12) and the identity (18) allows to speak about the existence of the structure on the space $A = A_2^1 \oplus A_1^2$ which is analogous to the Lie algebra on the matrix. So there is the question about the analogues of Lie groups and defining the exponential function in this case, if we start from the notion of automorphism as the action which preserve the structure. Note that 3-ary operation (12) and identity (18) can be defined on the pairs of rectangle matrices. It is possible that studying of this case is more simple then the case of tensors.

In the present paper we consider finite subalgebras of mixed tensor algebra $\bigoplus_{i,j=0}^{\infty} \mathcal{A}_i^j$. Note, that in [4] there were considered infinite subalgebras and their properties with respect to the nilpotent boundary operators with $\delta^k = 0$, $k \geq 3$. Another important class of questions arises from the point of view when the mixed tensor algebra is regarded as the one generated by a pair linear spaces with bilinear form. This assumes the generalization in which the free "algebras" are generated by n -tuples of spaces contracted via n -linear forms.

7 Appendix 1.

Let us consider the operation (2) with values $\alpha = +1$, $\beta = -1$, $\gamma = +1$, $\delta = -1$. We get:

$$A \circ B = A \cdot B - B \cdot A + A \cdot Tr B - B \cdot Tr A. \quad (20)$$

Show that $(A \circ B) \circ C + (C \circ A) \circ B + (B \circ C) \circ A \neq 0$. For this let us calculate:

$$\begin{aligned}
& (A \circ B) \circ C = \\
& = +ABC - BAC - CAB + CBA - TrC \cdot TrA \cdot B + TrC \cdot TrB \cdot A - \\
& - TrA \cdot BC + TrA \cdot CB + TrB \cdot AC - TrB \cdot CA + TrC \cdot AB - TrC \cdot BA \\
& (C \circ A) \circ B = \\
& = +CAB - ACB - BCA + BAC - TrB \cdot TrC \cdot A + TrB \cdot TrA \cdot C - \\
& - TrC \cdot AB + TrC \cdot BA + TrA \cdot CB - TrA \cdot BC + TrB \cdot CA - TrB \cdot AC - \\
& - TrB \cdot TrC \cdot A + TrB \cdot TrA \cdot C \\
& (B \circ C) \circ A = \\
& = +BCA - CBA - ABC + ACB - TrA \cdot TrB \cdot C + TrA \cdot TrC \cdot B - \\
& - TrB \cdot CA + TrB \cdot AC + TrC \cdot BA - TrC \cdot AB + TrA \cdot BC - TrA \cdot CB
\end{aligned} \tag{21}$$

Summing we have the following expression:

$$\begin{aligned}
& (A \circ B) \circ C + (C \circ A) \circ B + (B \circ C) \circ A = \\
& = TrA \cdot (CB - BC) + TrB \cdot (AC - CA) + TrC \cdot (BA - AB).
\end{aligned} \tag{22}$$

Now let us consider the result of the following expression $((A \circ B) \circ C + (C \circ A) \circ B + (B \circ C) \circ A) \circ D$:

$$\begin{aligned}
& ((A \circ B) \circ C + (C \circ A) \circ B + (B \circ C) \circ A) \circ D = \\
& = -TrA \cdot (BC - CB) \cdot D + TrB \cdot (CA - AC) \cdot D - TrC \cdot (AB - BA) \cdot D - \\
& + D \cdot TrA \cdot (BC - CB) + D \cdot TrB \cdot (CA - AC) + D \cdot TrC \cdot (AB - BA) - \\
& - TrD \cdot TrA \cdot (BC - CB) - TrD \cdot TrB \cdot (CA - AC) - TrD \cdot TrC \cdot (AB - BA).
\end{aligned} \tag{23}$$

Summing this result with the results of the other expressions:

$$\begin{aligned}
& ((C \circ B) \circ D + ((D \circ C) \circ B + (B \circ D) \circ C) \circ A, \\
& ((C \circ D) \circ A + (A \circ C) \circ D + (D \circ A) \circ C) \circ B, \\
& ((B \circ A) \circ D + (D \cdot B) \circ A + (A \circ D) \circ B) \circ C,
\end{aligned} \tag{24}$$

we get the value equal to zero - (6).

8 Appendix 2.

Each term in (16) gives $36 \cdot 2$ words. The number of terms is 20, so the number of different words is $720 \cdot 2$. There are the words distinguished by the belonging to the (2,1)-type or (1,2)-type tensors. So any word of each tensor type occurs 6 times because the total number of different words is $5! = 120$. Note that it is possible to compose 12 different words using the different symbols, for example: $A_1^2, B_2^1, C_1^2, D_2^1, E_1^2$. It means that the set having 120 different words divides in 10 classes. In each five-symbolic word there are alternating tensors belonging to the different types ((2,1) and (1,2)). So we can code the tensor type by the tensor type of second and fourth symbol. At the other hand there are 10 sets which consist of two symbol

chosen from given five ones. It defines the dividing of the set having 120 words of each tensor type in 10 classes consisted of 12 words.

Now let us consider any class, for example $\{a, e\}$. We can form the table in which to each word (left column) we associate the primary operations (upper levels). It help us to find the term of identity which participates in generating of considering word. In lower levels there are the weights after two operations:

A, E	abc, cba	ace, eca	cde, edc	bda, adb	aed, dea	ebd, dba	dac, cad	eab, bae	ceb, bec	
$baced$	+	+	+							$Eq1$
$badec$			+	+	+					$Eq2$
$bedac$					+	+	+			$Eq3$
$becad$		+					+		+	$Eq4$
$cabed$	+					+		+		$Eq2$
$cadeb$					+	+	+			$Eq3$
$cebad$				+				+	+	$Eq3$
$cedab$			+	+	+					$Eq2$
$dabec$				+				+	+	$Eq3$
$daceb$		+					+		+	$Eq4$
$decab$	+	+	+							$Eq1$
$debac$	+					+		+		$Eq2$
	$\beta\gamma, \gamma\alpha$	$\alpha\beta, \alpha\beta$	$\beta\gamma, \gamma\alpha$	$\gamma\gamma, \alpha\beta$	$\beta\beta, \beta\gamma$	$\gamma\gamma, \alpha\beta$	$\alpha\alpha, \alpha\gamma$	$\beta\beta, \beta\gamma$	$\alpha\alpha, \alpha\gamma$	

(25)

where:

$$\begin{aligned}
Eq1 : 2 \cdot (\beta\gamma + \gamma\alpha + \alpha\beta) &= 0 \\
Eq2 : \gamma\gamma + \gamma\alpha + \gamma\beta + \beta\beta + \beta\alpha + \beta\gamma &= 0 \\
Eq3 : \gamma\gamma + \beta\beta + \alpha\alpha + \gamma\alpha + \gamma\beta + \beta\alpha &= 0 \\
Eq4 : 2 \cdot (\alpha\alpha + \alpha\beta + \alpha\gamma) &= 0.
\end{aligned}
\tag{26}$$

So, the total weight of each word (after summing) is equal to zero. Note that this structure realizes in the other classes parameterized by two symbols.

Acknowledgments Authors thank G.Sharigin for the useful remarks and A.Morozov for fruitful discussions and attention to the work. The work of Yu. Chernyakov was supported in part by grant RFFI 03-02-17554 and grant 1999.2003.2 for support of scientific schools. The work of V.Dolotin was supported by the grant RFFI 04-02-17227.

References

- [1] Van der Varden B.L., Algebra, M., Nauka, 1976.
- [2] Shafarevich I.R., Osnovnie ponyatiya algebri. Sovremennye problemi matematiki. Fundamentalnye napravleniya, v.11, (VINITI), 1986.
- [3] D.Bar-Natan, S.Garoufalidis, L.Rozansky, D.P.Thurston, "The Aarhus integral of rational homology 3-spheres II: invariance and universality", qa/9801049.
- [4] V.Dolotin, "On invariant theory", alg-geom/9512011.